

Functional Somatic Syndromes

Arthur J. Barsky, MD, and Jonathan F. Borus, MD

The term *functional somatic syndrome* has been applied to several related syndromes characterized more by symptoms, suffering, and disability than by consistently demonstrable tissue abnormality. These syndromes include multiple chemical sensitivity, the sick building syndrome, repetition stress injury, the side effects of silicone breast implants, the Gulf War syndrome, chronic whiplash, the chronic fatigue syndrome, the irritable bowel syndrome, and fibromyalgia. Patients with functional somatic syndromes have explicit and highly elaborated self-diagnoses, and their symptoms are often refractory to reassurance, explanation, and standard treatment of symptoms. They share similar phenomenologies, high rates of co-occurrence, similar epidemiologic characteristics, and higher-than-expected prevalences of psychiatric comorbidity. Although discrete pathophysiologic causes may ultimately be found in some patients with functional somatic syndromes, the suffering of these patients is exacerbated by a self-perpetuating, self-validating cycle in which common, endemic, somatic symptoms are incorrectly attributed to serious abnormality, reinforcing the patient's belief that he or she has a serious disease. Four psychosocial factors propel this cycle of symptom amplification: the belief that one has a serious disease; the expectation that one's condition is likely to worsen; the "sick role," including the effects of litigation and compensation; and the alarming portrayal of the condition as catastrophic and disabling. The climate surrounding functional somatic syndromes includes sensationalized media coverage, profound suspicion of medical expertise and physicians, the mobilization of parties with a vested self-interest in the status of functional somatic syndromes, litigation, and a clinical approach that overemphasizes the biomedical and ignores psychosocial factors. All of these influences exacerbate and perpetuate the somatic distress of patients with functional somatic syndromes, heighten their fears and pessimistic expectations, prolong their disability, and reinforce their sick role. A six-step strategy for helping patients with functional somatic syndromes is presented here.

The term *functional somatic syndrome* refers to several related syndromes that are characterized more by symptoms, suffering, and disability than by disease-specific, demonstrable abnormalities of structure or function. Physicians in many medical specialties are increasingly confronted by patients who have disabling, medically unexplained, somatic symptoms and who have already arrived at a diagnostic label for their illness. The functional somatic syndromes have acquired major sociocultural and political dimensions. Their definitive status in public consciousness and popular discourse contrasts markedly with their still uncertain scientific and biomedical status. Patients with these syndromes often have very explicit disease attributions for their symptoms, and they resist information that contradicts these attributions (1, 2). These patients often have a strong sense of assertiveness and embattled advocacy with respect to their etiologic suppositions, and they may devalue and dismiss medical authority and epidemiologic evidence that conflicts with their beliefs (3).

The functional somatic syndromes include multiple chemical sensitivity, the sick building syndrome, repetition stress injury, chronic whiplash, chronic Lyme disease, the side effects of silicone breast implants, candidiasis hypersensitivity, the Gulf War syndrome, food allergies, mitral valve prolapse, and hypoglycemia. The incidence of several other functional somatic syndromes has apparently declined: chronic carbon monoxide poisoning; chronic mononucleosis; and symptoms resulting from exposure to video display terminals, carbonless copy paper, and weak electromagnetic fields. In three other syndromes—the chronic fatigue syndrome, fibromyalgia, and the irritable bowel syndrome—more uncertainty exists about the presence of demonstrable pathophysiology, but these syndromes are included in this review because they have extensive phenomenologic overlap with other functional somatic syndromes and the psychosocial factors discussed here apply to them.

This paper is also available at <http://www.acponline.org>.

Ann Intern Med. 1999;130:910-921.

From Brigham and Women's Hospital, Boston, Massachusetts. For current author addresses, see end of text.

Methods

English-language articles were identified through a search of the MEDLINE database from 1966 to

the present. The bibliographies of the retrieved articles were then searched for additional publications. Standardized or structured analysis of the identified papers was not possible because of variation in quality, design, and methods and because of the breadth of the articles included. Emphasis was given to empirical studies that used more rigorous diagnostic methods, larger samples, systematic analyses, appropriate comparison groups, and longitudinal follow-up.

The funding source for this work had no role in the collection, analysis, or interpretation of data or in the decision to submit the paper for publication.

Historical Context

In the past, various conditions associated with the symptoms of functional somatic syndromes (such as headache, musculoskeletal pain, fatigue, gastrointestinal distress, memory difficulties, and insomnia) have arisen, attracted intense medical attention, and then declined in incidence. Neurasthenia, spinal irritation, chronic brucellosis, pinched nerves, railway spine, and soldier's heart were each initially thought to have a medical cause, but when no pathologic basis for these conditions could be established, they subsequently declined in incidence and prevalence. More recently, functional somatic syndromes such as mercury poisoning caused by dental fillings, symptoms resulting from use of video display terminals, and chronic mononucleosis have declined in popularity.

Somatic distress and medically unexplained symptoms have always been endemic to daily life, but the social and cultural characteristics of each era shape the expression, interpretation, and attribution of these symptoms. Thus, similar constellations of benign symptoms acquire different diagnostic labels and are attributed to different causes in different time periods (1, 3). A line of descent can be traced from the DaCosta syndrome through soldier's heart, shell shock, and battle fatigue to the Gulf War syndrome (4). Musculoskeletal pain in the workplace, which previously manifested as writer's cramp and telegraphist's wrist, is now termed *repetition strain injury* (5). There are similarities between railway spine, common in the early 20th century, and the more recent chronic whiplash syndrome (6).

Although the functional somatic syndromes are not new, patients who have these syndromes today differ from their predecessors by being less relieved by negative findings on medical evaluation and less responsive to explanation, reassurance, and palliative treatment (1, 7, 8). Several factors may account for this shift.

First, the authority and prestige of the physician have declined: The reassurance of one's personal

physician and the opinions of medical and public health authorities are no longer as calming, reassuring, and palliative as they once were. With this erosion of physician authority and the increasing prevalence of a generalized antiscientific attitude (7), the determination that a functional somatic syndrome has no pathologic basis does not result in a rapid decline in the incidence of that syndrome, as it did in the past (8). This divergence of medical and scientific evidence and public opinion is particularly evident in the recent controversy over silicone breast implants (7).

Second, the current situation is powerfully shaped by the mass media (7–9), which often use hyperbole and uncritical reporting to portray the functional somatic syndromes (3, 7, 10, 11). Preliminary data, tentative findings, and the personal accounts of individual sufferers are reported as conclusive medical evidence (3, 12). The functional somatic syndromes are described as rapidly spreading epidemics, progressive and incapacitating, and some reports insinuate that powerful societal institutions are denying the existence of these syndromes to conceal their own negligence or culpability (3, 11). Such sensationalism and alarmism promote symptoms and distress (5, 13–17).

Finally, the contemporary climate is marked by the prominent political, legal, economic, and regulatory ramifications of the functional somatic syndromes (18–20). Individuals and organizations have strong vested interests in the status of these syndromes, and the actions of these persons and groups may reinforce sufferers' beliefs that their symptoms have a medical basis (21–24). The functional somatic syndromes form the basis for lawsuits and class actions seeking to attribute liability and fault. Medical specialists and clinics develop professional and financial stakes in one syndrome or another. Advocacy groups emerge to mobilize public opinion, influence scientific debate, and shape public policy. The functional somatic syndromes are a source of disputes over health insurance coverage; may propel the creation of environmental, occupational, and workplace regulations; and may qualify sufferers for worker's compensation or disability benefits.

Overlap and Common Characteristics

Each functional somatic syndrome is seen in a heterogeneous group of patients. In some patients, symptoms are attributable to a known disease entity; in others, they result from an unrecognized disorder that may involve physiologic or immunologic hyperreactivity and perceptual hypersensitivity. Other patients have symptoms that are caused by a

psychiatric disorder, and still others have symptoms that are best understood as a response to stressful life circumstances. Our knowledge of the functional somatic syndromes is incomplete, and we do not fully understand the etiologic roles of biological, psychological, and sociocultural factors in these syndromes. Although complex, poorly understood, and heterogeneous, the functional somatic syndromes nonetheless have enough in common to justify our discussing them together as variants of a common biopsychosocial process.

The similarities seen in the functional somatic syndromes have led some to propose that they share a common pathophysiology. Thus, they have been conceptualized as variants of “affective spectrum disorder” because a significant fraction of patients who have these syndromes respond to antidepressant medications of different, unrelated chemical classes (25–27). It has also been suggested that the functional somatic syndromes all involve the same pathophysiologic dysregulation and blunting of the central nervous system’s response to stress (28). Further research may shed light on these interesting hypotheses, but they are currently largely speculative.

Phenomenology

Although individual functional somatic syndromes may present with some organ-specific symptoms and may differ with respect to which symptoms are most prominent (for example, neck pain in chronic whiplash and gastrointestinal symptoms in the irritable bowel syndrome), they generally lack characteristic clinical presentations or distinct symptom complexes that are consistent across cases and that distinguish the syndromes from one other (29, 30). The various functional somatic syndromes have remarkably similar symptoms that share two important characteristics: They are diffuse, nonspecific, and ambiguous, and they are very prevalent in healthy, nonpatient populations (31). Symptoms common to the functional somatic syndromes include fatigue; weakness; sleep difficulties; headache; muscle aches and joint pain; problems with memory, attention, and concentration; nausea and other gastrointestinal symptoms; anxiety; depression; irritability; palpitations and “racing heart”; shortness of breath; dizziness or light-headedness; sore throat; and dry mouth.

All of these symptoms have a high incidence in the general population. Surveys of healthy persons who are not patients show that fatigue, headache, joint aches and stiffness, upper respiratory symptoms, and diarrhea are common and generally resolve spontaneously, usually within 1 month (32). Significant fatigue, for example, is reported by more than 20% of adults (33–36), and 30% of persons report current musculoskeletal symptoms (37).

Eighty-six percent to 95% of the general population has at least one somatic symptom in a given 2- to 4-week period (32, 38–41). The typical adult has a symptom every 4 to 6 days (32, 39), and 81% of healthy college students report having at least one somatic symptom in a 3-day period (42).

Overlap and Co-occurrence

The functional somatic syndromes have a high degree of overlap and co-occurrence (28, 43–45). Because the conditions are phenomenologically similar (31), the same person often meets the diagnostic criteria for several functional somatic syndromes simultaneously. Considerable overlap has been reported between multiple chemical sensitivity and repetition stress injury (46); between fibromyalgia and the chronic fatigue syndrome; between fibromyalgia and the irritable bowel syndrome (47–51); among multiple chemical sensitivity, the irritable bowel syndrome, and the Gulf War syndrome (31, 45, 50, 52, 53); and among the chronic fatigue syndrome, multiple chemical sensitivity, and fibromyalgia (31, 52). Over time, the same person may believe that he or she has several different functional somatic syndromes (43), a process referred to as *pathoplasticity* (1), and the diagnostic label given to a particular patient may be as strongly influenced by the context and medical specialty of the diagnostician as by the patient’s symptoms (54). Confronted with the same polysymptomatic patient, a rheumatologist may focus on upper-extremity symptoms and diagnose repetition stress injury, an internist may inquire into constitutional symptoms and suspect the chronic fatigue syndrome, an allergist may diagnose the sick building syndrome, and a gastroenterologist may focus on bowel symptoms and identify the irritable bowel syndrome (54).

Epidemiology

The functional somatic syndromes have several epidemiologic similarities. They often begin in limited, sporadic outbreaks among small groups of people who are in close contact with each other (such as residents of small, rural towns; coworkers in the same office; or members of particular military units) and then “spread” to other persons with similar risk profiles after widespread publicity and alarm (1–3, 46, 55–60). The pattern of these epidemic-like outbreaks at first suggests infectious contagion or a common toxic agent. Epidemiologic scrutiny, however, shows that the spread occurs along lines of interpersonal communication, acquaintance, and familiarity as well as with physical proximity or exposure to the suspected “pathogen” (1, 5, 18, 20, 43, 46, 61). Prevalence rates vary widely in similar populations exposed to the same putative etiologic agent (62–64), and similar groups in various geo-

graphic locations do not necessarily develop a given syndrome (for example, keyboard operators outside of the United States, England, and Australia develop repetition stress injury less frequently, and non-English-speaking troops deployed to the Persian Gulf did not develop the Gulf War syndrome as often). No dose-response relation can be firmly established (5, 16, 66–70), and no pathogenic toxin, infectious agent, or physical vector is discovered after extensive evaluation (5, 46, 68, 71).

Comorbid Psychiatric Disorders

Patients with functional somatic syndromes have elevated rates of psychiatric disorders, particularly anxiety, depressive, and somatoform disorders. The cause-and-effect relation between the functional somatic syndromes and psychiatric disorders is widely debated because it is often difficult to determine which condition is antecedent and which is consequent (72). Nonetheless, the prevalence of axis I psychiatric disorders, both current and lifetime, is clearly higher in patients with functional somatic syndromes than in the general population or in similar groups of medically ill patients (18–20, 28, 44, 49, 50, 71–75). For example, the prevalence of psychiatric symptoms and psychiatric diagnoses is significantly higher in patients with fibromyalgia than in patients with rheumatoid arthritis or in healthy persons (25, 26, 76–80). Patients with the irritable bowel syndrome have more psychiatric diagnoses, personality disorders, and psychiatric symptoms than patients with inflammatory bowel disease do (81–84). The prevalence of premorbid and current psychiatric disorders is higher in patients with multiple chemical sensitivity than in numerous comparison groups (18, 19, 29, 71, 85), and elevated rates of anxiety and depressive disorders have been seen in several populations with the chronic fatigue syndrome (44, 73, 86–93).

Patients with functional somatic syndromes, including those with the chronic fatigue syndrome, multiple chemical sensitivity, fibromyalgia, and the irritable bowel syndrome, also have a higher prevalence of somatization, somatoform disorders, and medically unexplained symptoms that are unrelated to the functional somatic syndromes (19, 20, 44, 71, 72, 84, 92, 94–101). In some studies, somatization (the experiencing of somatic symptoms that do not have a demonstrable medical basis, the belief that these symptoms are due to demonstrable disease, and the seeking of medical attention for them) predates the onset of the functional somatic syndromes (4, 74); this suggests a preexisting tendency to experience and report bodily distress. For example, a group of patients with multiple chemical sensitivity had significantly more medically unexplained somatic symptoms and a higher prevalence of soma-

tization disorder before the onset of multiple chemical sensitivity than a comparison group did (71). However, somatization occurs in almost everyone at some time and to some degree and does not itself indicate a psychiatric disorder. Because the functional somatic syndromes are determined by multiple factors and are much shaped by psychological, sociocultural, and circumstantial forces, they resist localization anywhere within our medical or psychiatric taxonomy.

Refractoriness to Treatment of Symptoms

The functional somatic syndromes are often refractory to usual medical treatments and standard palliative measures (2). Epidemiologic comparisons of patients who have self-diagnosed functional somatic syndromes with community residents who report the same symptoms suggest that refractoriness, chronicity, and intractability of symptoms are more characteristic of the former group. In those functional somatic syndromes for which an environmental cause is postulated, improvement does not reliably result from control or elimination of the putative toxic agent (46, 102–104). When a physical activity is thought to be pathogenic, rest and physiotherapy are generally not effective (46, 104–106). When restriction of a patient's activities and functioning fails to relieve a given symptom, this is often regarded not as evidence against the putative cause-and-effect relation but rather as an indication that the restrictions were not stringent enough. Patients are thus caught in a vicious cycle in which the ineffectiveness of a treatment strategy leads to its intensification rather than its abandonment.

Amplification and Maintenance of Somatic Symptoms

An Explanatory Model

No single mechanism accounts for the functional somatic syndromes, but the knowledge we have is enough to suggest an explanatory model for the genesis and maintenance of these conditions (11, 107). Distressing symptoms are omnipresent in daily life. They result from benign dysfunctions and self-limited ailments; chronic medical conditions; psychosocial stress; psychiatric disorders; and, less frequently, previously unknown or unrecognized medical conditions. Under the influence of medical scrutiny, public health concern, and media attention, a process of symptom amplification that alters the perception of these endemic symptoms can be set in motion. Learning about a disease of which we were previously unaware (through personal contact with a sufferer, word of mouth, or the media) may lead us

to tentatively reattribute previously ill-defined or treatment-resistant chronic symptoms to the “new” disease (74, 108). (For example, nasal stuffiness and headaches may be ascribed to the sick building syndrome.) This reattribution then amplifies the symptoms themselves, making them seem more intense, noxious, and troublesome (107, 109, 110). The assumption that one is seriously ill also heightens self-scrutiny and prompts a confirmatory search for other symptoms to corroborate one’s suspicions. Ambiguous sensations that were previously ignored, dismissed as innocuous, or never consciously noticed are now interpreted as further evidence of the presence of the suspected disease (107, 109, 110). A self-validating and self-perpetuating cycle of symptom amplification and disease conviction ensues: The suspicion of disease heightens bodily awareness, symptom perception, and distress, and these, in turn, reinforce the belief that the sufferer is sick.

This process of confirmatory bias and symptom amplification operates in each individual sufferer. It may also serve as a mechanism for “transmitting” the syndrome from one person to another. A new syndrome may first appear when a few persons with an unusual or previously unknown or ill-defined medical condition are recognized. Under the influence of growing medical and public attention, these persons serve as a nidus around which aggregate other persons who have similar symptoms but do not actually have the same underlying condition. Media publicity, sympathetic physicians, special clinics devoted to the condition, hotlines, litigation, disability compensation, and patient advocacy groups serve as vectors and propel this amplification of symptoms and reattribution of preexisting somatic distress. This process is mediated by four mechanisms: the belief that one has a disease, negative expectations about the future course of the disease, the sick role, and stressful events.

The few persons originally affected may serve as a template for others with similar, preexisting symptoms who reattribute their symptoms to the functional somatic syndrome about which they have recently learned. Sociocultural forces then reinforce the reattribution and, ultimately, the symptoms themselves. Some persons (for example, those with a history of trauma, those with psychiatric disorders, those undergoing major life stress, and those whose families or caregivers reinforce their symptoms and illness behavior [2, 111, 112]) are more vulnerable to this process of amplification. The following discussion focuses on the four specific mechanisms involved in symptom amplification. These amplifiers were selected because they are particularly salient in perpetuating and in maintaining patient distress and because we have empirical evidence about their roles.

Psychosocial Factors That Amplify Symptoms

The Belief That One Is Sick

Bodily perception is an active, not passive, process. Myriad somatic and visceral stimuli are constantly filtered in the brain, and only a small fraction reach conscious attention (109, 113–115). Our suspicions about the causes of our sensations guide this filtering and appraisal process: Sensations thought to have pathologic significance are selected for conscious attention and are amplified. The influence of cognitive beliefs on somatic perception is evident in studies showing that disease labeling results in decreased psychological health and increased absenteeism (116). For example, patients who did not know that they were hypertensive show a threefold increase in days of work missed after diagnosis; this effect is independent of the anti-hypertensive regimen (117). In a prospective study of herpes zoster (118), the persistence of pain at follow-up was predicted by the extent of the patient’s conviction about the disease at inception. Among patients with chest pain but not serious coronary artery disease, the persistence of pain was predicted by the patient’s earlier belief that he or she was prone to serious heart disease (119). Similarly, the persistence of fatigue after viral infection has been associated with the patient’s belief in his or her vulnerability to viruses and with the tendency to ascribe ambiguous bodily symptoms to disease (120). In two prospective studies of the chronic fatigue syndrome (22, 23), the strength of the sufferers’ belief that their fatigue had a medical basis predicted poor subsequent symptomatic outcome. Finally, patients’ convictions that they had severe lactose intolerance led them to misattribute various benign abdominal symptoms to this disorder (67).

Beliefs about disease also bias recall of past symptoms (121). In a comparison with uninjured controls, patients with whiplash were found to underestimate their preinjury history of neck symptoms (62, 122). Compared with women who had less negative views of menstrual distress, women who believed that menstruation is a negative experience recalled past menstrual periods as more symptomatic than they had reported them to be when they were experiencing them. The two groups of women did not differ, however, in their recall of intermenstrual symptoms (123). Similarly, informing healthy volunteers in an experiment that they had just tested positive for a disease caused them to recall symptoms that were said to characterize that disease and to recall more behaviors that were described as risk factors for the disease (124).

Thus, the more convinced patients with functional somatic syndromes are that their symptoms are serious and pathologic, the more intense, pro-

longed, and disabling the symptoms become. Such symptom amplification is fostered by physicians who prematurely focus exclusively on medical explanations for the symptoms, by alarming anecdotes in the popular press and on the Internet, and by organized campaigns to designate a particular syndrome as a serious disease (24, 125, 126).

Future Expectations and the Role of Suggestion

Suggestion amplifies and maintains symptoms because humans tend to perceive what they expect to perceive. The cognitive processing of current bodily sensation is guided by our expectations of what we will experience next. This was shown in a multicenter study of aspirin treatment for unstable angina (127). Patients whose informed consent forms explicitly mentioned possible gastrointestinal side effects had a significantly higher incidence of gastrointestinal symptoms (but not confirmed gastrointestinal disease) than did patients whose forms did not specifically mention these effects. Six times as many patients in the former group withdrew from the study because of gastrointestinal distress (127). In patients with the chronic fatigue syndrome, extremely negative expectations about the future consequences of exercise are associated with higher levels of fatigue and disability (128). Similarly, patients who are more concerned about the seriousness of whiplash at the time of injury have longer lasting symptoms (129), and the expectations of patients with mild head injury with regard to future symptoms explain as much of the variance in symptoms as the injuries themselves (62). The power of suggestion has also been shown to influence healthy persons: Instructing persons to attend to evidence of "nasal obstruction" as they breathed induced more symptoms than instructing them to attend to the "free passage of air" (109). Similarly, headache was induced in volunteers who were told that a mild electric current that produces headache would be passed through their heads, when in fact no electricity was administered (130).

Studies of communities exposed to toxic waste pollution are also relevant. Compared with residents of unexposed communities, exposed residents report a broader range of somatic symptoms than can be attributed to the pollutant, and symptoms are most prominent in persons who believe that toxic waste and environmental pollution are more threatening and dangerous (131–133). Media coverage, community and legal action, and allegations of cover-ups alter the perception of normally occurring benign symptoms in those who expect to become sick, causing them to misattribute symptoms to the pollutant (131, 132). One explanation for the increased incidence of somatic symptoms in Gulf War veterans may be the suggestions made by the media, some

medical professionals, and advocacy groups about the negative health consequences of suspected toxic exposures (126). Similarly, persons investigating repetition stress injury have concluded that exaggerated media reports of this condition's seriousness and suggestions that the condition is progressive and incapacitating perpetuate the symptoms and disability associated with it (5, 16).

The Sick Role

Symptoms are also amplified by the act of becoming a patient. The assumption of the sick role can initiate far-reaching and pervasive changes—such as unemployment, altered social relationships and family dynamics, and medical help seeking—that in themselves amplify symptoms. Thus, the responses of family members, employers, and physicians to a patient's illness behavior can exacerbate or alleviate chronic pain and the symptoms of somatoform disorders (134–137), and the chronicity of medically unexplained symptoms has been empirically associated with such "secondary gains" (138). In general terms, social labeling theory posits that the connotations and implications of the label we apply to a condition or state influence the outcome of that condition or state. Once a person is labeled as ill, for example, he or she is regarded and treated in ways that make recovery more difficult: Continued illness is expected of the person, and symptoms therefore persist (139).

Health-contingent litigation, monetary compensation, and disability payments all have negative effects on symptoms (140). This was shown by a recent study of whiplash in a country that has little physician or public awareness of the syndrome, no litigation or compensation for it, and no involvement of insurance companies. Victims of rear-end motor vehicle accidents in this country did not have a higher incidence of postaccident headache and neck pain than did randomly chosen, uninjured, age- and sex-matched persons (63). A large body of literature indicates that injury compensation and worker disability payments are associated with a poorer symptomatic outcome after medical treatment and with a prolonged rehabilitative course (141–143). For example, recovery from surgery for the carpal tunnel syndrome is more prolonged and more symptomatic in persons who receive workers' compensation than in those who do not (144). Forcing someone to repeatedly prove that he or she is sick confounds the illness experience, impedes recovery from symptoms, and fosters invalidism (142, 145). When the continuation of benefits is contingent on the continuation of symptoms, the patient is trapped in the sick role. Thus, the incidence of repetitive stress injury is closely correlated with the availability and generosity of disability and workers'

compensation payments, and it declines after administrative and judicial decrees that it is not compensable (16, 104, 146).

When persons with functional somatic syndromes become patients and are given a diagnosis, they are admitted to the sick role. They may curtail or stop work, limit recreational or social activities, pursue legal action or receive disability compensation; read about their condition in magazines and on the Internet, meet and talk with fellow sufferers, and join an advocacy group. Although these steps may be adaptive and appropriate for some, they may also have unintended, long-term, negative consequences by strengthening expectations of future distress, reinforcing symptoms, and making recovery more difficult. Recovery is more difficult and requires greater face saving when sick role behaviors have become more extensive and ingrained: Clinical improvement may seem to call into question the patient's veracity or the legitimacy of his or her symptoms.

Stress and Distress

Stress exacerbates and perpetuates physical symptoms, lowers the threshold for medical help seeking, and makes us quicker to conclude that an ambiguous bodily sensation is due to disease (111, 147–152). Two types of stress are relevant: 1) daily life problems and recurring minor irritants and 2) major life changes and events requiring adaptation. Repetition stress injury, for example, is closely associated with daily stresses and hassles in the workplace and tends to occur when workers must adapt to a new technology that is demanding, threatens job security, and raises expectations for productivity. Clerical workers who report upper-extremity pain also report greater work demands, less control over their work, more job insecurity, and less camaraderie with their coworkers than do workers without such pain (153, 154). A similar relation exists between job stress and back pain (155–158), and perceived work intensity, mental strain, and stressful home lives are more common among workers who acutely develop the sick building syndrome (159). Recurrent, daily stresses have been shown to amplify pain in patients with rheumatoid arthritis (160, 161). Similarly, chronic whiplash symptoms 6 months after a motor vehicle accident were prospectively predicted by daily life stresses in the months before the injury, whereas neurologic signs did not predict subsequent distress (66, 162).

Stressful major life events have also been shown to amplify bodily symptoms. Natural disasters, such as floods (163–165); warfare; criminal victimization; and exposure to environmental pollutants (17, 166–168) result in medically unexplained symptoms. Emotionally laden stressors have been found to exacerbate or precipitate many functional somatic syn-

dromes (28, 169–171). Military combat has resulted in a consistent syndrome of medically unexplained symptoms in U.S. soldiers since the U.S. Civil War (4, 17, 172). Medically unexplained somatic symptoms increase substantially in populations stressed by exposure to environmental toxins and in populations that are only rumored to have had such exposure (13–15).

Stress amplifies symptoms in two ways. First, because stress is widely known to be pathogenic, persons under stress are quicker to ascribe ambiguous bodily symptoms to disease rather than to attribute them to normal physiology, as they might otherwise do. Second, external stressors induce anxiety and depression, which have their own somatic and autonomic concomitants. Anxiety decreases the pain threshold and pain tolerance (173). It also causes apprehensive self-scrutiny and a sense of physical threat and jeopardy, which make symptoms more noxious, ominous, and worrisome (174–176). Depression, in addition to producing its own autonomic symptoms, amplifies and perpetuates other somatic symptoms (174–176). For example, patients who had more persistent and prolonged symptoms after an influenza outbreak were shown to have had higher levels of depression before becoming sick (177).

The more the functional somatic syndromes are thought of as ominous, incapacitating, and severe, and the more alarm and peril are associated with them (in short, the more stressful the experience of illness), the more intense and disabling symptoms become (5, 13–17).

Helping the Patient

The hyperbole, litigation, compensation, and self-interested advocacy surrounding the functional somatic syndromes can exacerbate and perpetuate symptoms, heighten fears and concerns, prolong disability, and reinforce the sick role. Excessive medical testing and treatment expose patients to iatrogenic harm and amplify symptoms. Exclusive emphasis on a search for structural abnormalities can distract physicians from eliciting the patient's beliefs, expectations, and personal circumstances. Patients with functional somatic syndromes can become so engrossed in establishing the legitimacy of their condition, so invested in discovering the cause of their symptoms, and so preoccupied with assigning fault and culpability that palliative treatment is made more difficult or is forgone.

Given these caveats, how should clinicians proceed? Medical management rests on six steps: 1) ruling out the presence of diagnosable medical disease, 2) searching for psychiatric disorders, 3) build-

ing a collaborative alliance with the patient, 4) making restoration of function the goal of treatment, 5) providing limited reassurance, and 6) prescribing cognitive-behavioral therapy for patients who have not responded to the aforementioned five steps.

First, clinicians must uphold their medical mandate with an appropriate search for a previously unrecognized medical disorder. In deciding how extensive this medical work-up should be, physicians must remember the adverse effects of overly aggressive investigation, of fostering the sick role, and of leading patients to expect a definitive medical explanation for all somatic distress. Caution is advised in ordering tests and obtaining specialty consultations solely to reassure the patient—negative findings provide little reassurance to most patients with chronic, medically unexplained symptoms and often ultimately heighten rather than assuage worry and anxiety (178–180). Furthermore, extensive medical testing carries the risk for iatrogenesis and solidifies the patient's conviction that his or her distress has a biomedical cause (24, 181). It is therefore helpful to have evidence-based guidelines for the appropriate extent of medical evaluation and the frequency with which such evaluation should be repeated. Currently, expert consensus has been promulgated for only a few functional somatic syndromes.

Second, the physician should search for diagnosable psychiatric disorders, particularly major depression and panic disorder (which are highly prevalent and treatable). Self-report screening questionnaires and brief, structural diagnostic interviews can assist the physician in this search. It is important to remember that the likelihood of a psychiatric diagnosis increases linearly with the number of somatic symptoms that the patient reports (97, 182–184). For example, compared with patients who have no pain, those who have medically unexplained pain at two sites have a fivefold higher prevalence of major depression, and those with three or more pains have eight times the risk for major depression (185). The stigma associated with a psychiatric diagnosis often makes patients feel that the legitimacy of their illness is being discounted and may make them cling more assiduously to a biomedical explanation of their symptoms (134). Patients must be assured that the presence of a psychiatric disorder in no way means that their somatic symptoms are imaginary or feigned. They should be told that psychiatric disorders are regarded less as causes of somatic symptoms than as amplifiers that exacerbate and perpetuate symptoms and impede recovery.

Third, a collaborative therapeutic alliance between physician and patient is crucial. The physician must take special care to acknowledge and legitimize the patient's suffering because a definitive biomedical explanation for the patient's symptoms has

proven elusive. At the same time, the physician should discourage the patient from assuming the sick role, should undercut alarming expectations about the clinical course, and should avoid making distressing symptom attributions. Closely related to the establishment of a collaborative alliance is the process of making symptom palliation, coping, and rehabilitation the focus of the clinical enterprise. The goal of treatment becomes the identification and alleviation of factors that amplify and perpetuate the patient's symptoms and cause functional impairment. The focus is on coping rather than on curing, on improving functional status rather than eradicating symptoms. If this is to be accomplished, patients with functional somatic syndromes must be actively involved in the treatment process and must be dissuaded from assuming a passive role and waiting to be cured by medical procedures or interventions. Realistic, incremental goals should be set and should be specified in terms of observable behaviors. (For example, a gently graduated exercise program should be prescribed.) Patients should be encouraged to resume their activities as much as possible and to remain at work if they are at all able.

Limited, cautious reassurance is appropriate. Patients can be reassured that grave medical diagnoses have been ruled out and can be told clearly that they do not have a lethal or progressive disease. However, because these patients feel ill and symptomatic, it is not enough to tell them what they do not have without telling them what they do have. It is often helpful to describe the process of amplification, whereby sociocultural and psychological processes exacerbate distress and hinder recovery. Although it does not provide a definitive etiologic explanation for a patient's distress, such a discussion gives patients an explanatory model that focuses on processes and functioning rather than on structural abnormalities.

Finally, if these strategies are insufficient, cognitive-behavioral therapies can be effective in treating the persistent distress and disability resulting from functional somatic syndromes. Such therapies have been developed for the somatoform disorders and for some medically unexplained symptoms, including those of the irritable bowel syndrome, fibromyalgia, the chronic fatigue syndrome, headache, and atypical chest pain (186–198). Controlled intervention trials with long-term follow-up have shown the effectiveness of cognitive-behavioral treatment in reducing somatic symptoms, generalized distress, and disability (186–197, 199–204). These interventions help patients cope with symptoms by helping them reexamine their health beliefs and expectations and explore the effects of the sick role and of stress and distress on their symptoms. They help patients find

alternative explanations for symptoms, restructure faulty disease beliefs, alter expectations, and learn techniques of focused attention and distraction. Behavioral strategies, such as response prevention, systematic desensitization, graduated exercise regimens, and progressive muscle relaxation, help those with functional somatic syndromes resume normal activities, minimize role impairment, and curtail sick role behaviors. The cognitive-behavioral approach stimulates patients to assume a more active role in coping and rehabilitation, and it counters the assumption that cure results only from the application of technological interventions to passive patients.

The role of traditional psychotherapy is generally restricted to cases in which the patient with a functional somatic syndrome identifies a psychological problem or a source of emotional distress for which he or she wants treatment. Psychotropic medications are indicated when a pharmacologically responsive psychiatric disorder (such as major depression or panic disorder) is present. In addition, antidepressants sometimes alleviate somatic symptoms (particularly pain and insomnia) and may improve the functional status of patients who have functional somatic syndromes and subthreshold psychiatric disorders. The empirical evidence for the efficacy of antidepressants is strongest for the chronic fatigue syndrome, fibromyalgia, and the irritable bowel syndrome (27). Little is known about the use of alternative therapies in functional somatic syndromes. They may help some patients by providing an enhanced sense of self-efficacy and control over symptoms, but empirical data on this topic are not available.

Conclusions

The functional somatic syndromes cause great suffering, distress, and disability and have substantial societal costs. The public, therefore, needs 1) to be cautioned about prematurely concluding that symptoms indicate serious disease, 2) to become more cognizant of the ubiquity of benign symptoms and self-limited conditions, and 3) to appreciate the influence of psychosocial factors on the experience of illness. Research into the functional somatic syndromes must continue, but the search for biological and physical causes of symptoms should be accompanied by study of psychological and sociocultural factors. Finally, the media must offer the public a less sensational, more accurate, and more sophisticated model of the functional somatic syndromes—one that encompasses both biomedical and psychosocial factors. Such a comprehensive, biopsychosocial approach to functional somatic syndromes by the medical profession, the public, and the media should

permit us to better understand and more effectively treat these conditions.

Acknowledgments: The authors thank Thomas Delbanco, MD, George E. Vaillant, MD, and Leon Eisenberg, MD, for assistance in preparing this manuscript.

Grant Support: In part by research grant MH-40487 from the National Institute of Mental Health (Bethesda, Maryland).

Requests for Reprints: Arthur J. Barsky, MD, Division of Psychiatry, Brigham and Women's Hospital, 75 Francis Street, Boston, MA 02115.

Current Author Addresses: Drs. Barsky and Borus: Division of Psychiatry, Brigham and Women's Hospital, 75 Francis Street, Boston, MA 02115.

References

1. **Shorter E.** From Paralysis to Fatigue: A History of Psychosomatic Illness in the Modern Era. New York: Free Pr; 1992.
2. **Kellner R.** Psychosomatic Syndromes and Somatic Symptoms. Washington, DC: American Psychiatric Pr; 1991.
3. **Showalter E.** Hysterics: Hysterical Epidemics and Modern Culture. New York: Columbia Univ Pr; 1997.
4. **Hyams KC, Wignall FS, Roswell R.** War syndromes and their evaluation: from the U.S. Civil War to the Persian Gulf War. *Ann Intern Med.* 1996;125:398-405.
5. **Ireland DC.** Psychological and physical aspects of occupational arm pain. *J Hand Surg [Br].* 1988;13:5-10.
6. **Trimble MR.** Post-Traumatic Neurosis: From Railway Spine to Whiplash. New York: J Wiley; 1981.
7. **Angell M.** Science on Trial: The Clash of Medical Evidence and the Law in the Breast Implant Case. New York: WW Norton; 1996.
8. **Shorter E.** Sucker-punched again! Physicians meet the disease-of-the-month syndrome. *J Psychosom Res.* 1995;39:115-8.
9. **Brown J, Chapman S, Lupton D.** Infinitesimal risk as a public health crisis: news media coverage of a doctor-patient HIV contact tracing investigation. *Soc Sci Med.* 1996;43:1685-95.
10. **Campion EW.** Power lines, cancer, and fear [Editorial]. *N Engl J Med.* 1997;337:44-6.
11. **Barsky AJ.** Worried Sick: Our Troubled Quest for Wellness. Boston: Little, Brown; 1988.
12. **Jauchem JR.** Epidemiologic studies of electric and magnetic fields and cancer: a case study of distortions by the media. *J Clin Epidemiol.* 1992;45:1137-42.
13. **Selden BS.** Adolescent epidemic hysteria presenting as a mass casualty, toxic exposure incident. *Ann Emerg Med.* 1989;18:892-5.
14. **Struwing JP, Gray GC.** An epidemic of respiratory complaints exacerbated by mass psychogenic illness in a military recruit population. *Am J Epidemiol.* 1990;132:1120-9.
15. **Small GW, Borus JF.** The influence of newspaper reports on outbreaks of mass hysteria. *Psychiatr Q.* 1987;58:269-78.
16. **Hall W, Morrow L.** "Repetition strain injury": an Australian epidemic of upper limb pain. *Soc Sci Med.* 1988;27:645-9.
17. **Decoufle P, Holmgren P, Boyle CA, Stroup NE.** Self-reported health status of Vietnam veterans in relation to perceived exposure to herbicides and combat. *Am J Epidemiol.* 1992;135:312-23.
18. **Black DW, Rathe A, Goldstein RB.** Environmental illness. A controlled study of 26 subjects with "20th century disease." *JAMA.* 1990;264:3166-70.
19. **Stewart DE, Raskin J.** Psychiatric assessment of patients with "20th-century disease" ("total allergy syndrome"). *Can Med Assoc J.* 1985;133:1001-6.
20. **Brodsky CM.** "Allergic to everything": a medical subculture. *Psychosomatics.* 1983;24:731-2, 734-6, 740-2.
21. **Black DW.** Iatrogenic (physician-induced) hypochondriasis. Four patient examples of "chemical sensitivity." *Psychosomatics.* 1996;37:390-3.
22. **Wilson A, Hickie I, Lloyd A, Hadzi-Pavlovic D, Boughton C, Dwyer J, et al.** Longitudinal study of outcome of chronic fatigue syndrome. *BMJ.* 1994;308:756-9.
23. **Vercoulen JH, Swanink CM, Fennis JF, Galama JM, van der Meer JW, Bleijenberg G.** Prognosis in chronic fatigue syndrome: a prospective study on the natural course. *J Neurol Neurosurg Psychiatr.* 1996;60:489-94.
24. **Kouyanou K, Pither CE, Wessely S.** Iatrogenic factors and chronic pain. *Psychosom Med.* 1997;59:597-604.
25. **Hudson JI, Pope HG Jr.** Fibromyalgia and psychopathology: is fibromyalgia a form of "affective spectrum disorder?" *J Rheumatol Suppl.* 1989;19:15-22.
26. **Hudson JI, Hudson MS, Pliner LF, Goldenberg DL, Pope HG Jr.** Fibromyalgia and major affective disorder: a controlled phenomenology and family history study. *Am J Psychiatry.* 1985;142:441-6.
27. **Gruber AJ, Hudson JI, Pope HG Jr.** The management of treatment-resistant depression in disorders on the interface of psychiatry and medicine.

- Fibromyalgia, chronic fatigue syndrome, migraine, irritable bowel syndrome, atypical facial pain, and premenstrual dysphoric disorder. *Psychiatr Clin North Am.* 1996;19:351-69.
28. **Clauw DJ, Chrousos GP.** Chronic pain and fatigue syndromes: overlapping clinical and neuroendocrine features and potential pathogenic mechanisms. *Neuroimmunomodulation.* 1997;4:134-53.
 29. **Terr AI.** Environmental illness. A clinical review of 50 cases. *Arch Intern Med.* 1986;146:145-9.
 30. **Hyams KC.** Lessons derived from evaluating Gulf War syndrome: suggested guidelines for investigating possible outbreaks of new diseases [Editorial]. *Psychosom Med.* 1998;60:137-9.
 31. **Buchwald D, Garrity D.** Comparison of patients with chronic fatigue syndrome, fibromyalgia, and multiple chemical sensitivities. *Arch Intern Med.* 1994;154:2049-53.
 32. **Verbrugge LM, Ascione FJ.** Exploring the iceberg. Common symptoms and how people care for them. *Med Care.* 1987;25:539-69.
 33. **Walker EA, Katon WJ, Jemelka RP.** Psychiatric disorders and medical care utilization among people in the general population who report fatigue. *J Gen Intern Med.* 1993;8:436-40.
 34. **Fukuda K, Dobbins JG, Wilson LJ, Dunn RA, Wilcox K, Smallwood D.** An epidemiologic study of fatigue with relevance for the chronic fatigue syndrome. *J Psychiatr Res.* 1997;31:19-29.
 35. **Pawlikowska T, Chalder T, Hirsch SR, Wallace P, Wright DJ, Wessely SC.** Population based study of fatigue and psychological distress. *BMJ.* 1994;308:763-6.
 36. **Buchwald D, Umali P, Umali J, Kith P, Pearlman T, Komaroff AL.** Chronic fatigue and the chronic fatigue syndrome: prevalence in a Pacific Northwest health care system. *Ann Intern Med.* 1995;123:81-8.
 37. **Cunningham LS, Kelsey JL.** Epidemiology of musculoskeletal impairments and associated disability. *Am J Public Health.* 1984;74:574-9.
 38. **Hannay DR.** The Symptom Iceberg: A Study of Community Health. Boston: Routledge & Kegan Paul; 1979.
 39. **Dunnell K, Cartwright A.** Medicine Takers, Prescribers, and Hoarders. London: Routledge & Kegan Paul; 1972.
 40. **Wadsworth ME, Butterfield WJ, Blaney R.** Health and Sickness: The Choice of Treatment, Perception of Illness and Use of Services in an Urban Community. London: Tavistock; 1971.
 41. **White KL, Williams TF, Greenberg BG.** The ecology of medical care. *N Engl J Med.* 1961;265:885-92.
 42. **Reidenberg MM, Lowenthal DT.** Adverse nondrug reactions. *N Engl J Med.* 1968;279:678-9.
 43. **Stewart DE.** The changing faces of somatization. *Psychosomatics.* 1990;31:153-8.
 44. **Manu P, Lane TJ, Matthews DA.** Somatization disorder in patients with chronic fatigue. *Psychosomatics.* 1989;30:388-95.
 45. **Wessely S.** Chronic fatigue and fibromyalgia syndromes. In: Sartorius N, ed. *Psychological Disorders in Medical Settings.* Berne: Hogrefe & Huber; 1990:82-97.
 46. **Göthe CJ, Molin C, Nilsson CG.** The environmental somatization syndrome. *Psychosomatics.* 1995;36:1-11.
 47. **Campbell SM, Clark S, Tindall EA, Forehand ME, Bennett RM.** Clinical characteristics of fibrositis. I. A "blinded," controlled study of symptoms and tender points. *Arthritis Rheum.* 1983;26:817-24.
 48. **Yunus M, Masi AT, Calabro JJ, Miller KA, Feigenbaum SL.** Primary fibromyalgia (fibrositis): clinical study of 50 patients with matched normal controls. *Semin Arthritis Rheum.* 1981;11:151-71.
 49. **Goldenberg DL.** Fibromyalgia syndrome. An emerging but controversial condition. *JAMA.* 1987;257:2782-7.
 50. **Hudson JI, Goldenberg DL, Pope HG Jr, Keck PE Jr, Schlesinger L.** Comorbidity of fibromyalgia with medical and psychiatric disorders. *Am J Med.* 1992;92:363-7.
 51. **Sandler RS, Drossman DA, Nathan HP, McKee DC.** Symptom complaints and health care seeking behavior in subjects with bowel dysfunction. *Gastroenterology.* 1984;87:314-8.
 52. **Goldenberg DL, Simms RW, Geiger A, Komaroff AL.** High frequency of fibromyalgia in patients with chronic fatigue seen in a primary care practice. *Arthritis Rheum.* 1990;33:381-7.
 53. **Miller CS.** Chemical sensitivity: history and phenomenology. *Toxicol Ind Health.* 1994;10:253-76.
 54. **Blackwell B.** Sick-role susceptibility. A commentary on the contemporary data base (1989-1991) and classification system. *Psychother Psychosom.* 1992;58:79-90.
 55. **Greenberg DB.** Neurasthenia in the 1980s: chronic mononucleosis, chronic fatigue syndrome, and anxiety and depressive disorders. *Psychosomatics.* 1990;31:129-37.
 56. **Small GW, Propper MW, Randolph ET, Eth S.** Mass hysteria among student performers: social relationship as a symptom predictor. *Am J Psychiatry.* 1991;148:1200-5.
 57. **Boxer PA.** Indoor air quality: a psychosocial perspective. *J Occup Med.* 1990;32:425-8.
 58. **Johanning E, Auger PL, Reijula K.** Building-related illnesses [Letter]. *N Engl J Med.* 1998;15:1070-1.
 59. **Chester AC, Levine PH.** The natural history of concurrent sick building syndrome and chronic fatigue syndrome. *J Psychiatr Res.* 1997;31:51-7.
 60. **Fukuda K, Nisenbaum R, Stewart G, Thompson WW, Robin L, Washko RM, et al.** Chronic multisymptom illness affecting Air Force veterans of the Gulf War. *JAMA.* 1999;280:981-8.
 61. **Levine PH, Jacobson S, Pocinki AG, Cheney P, Peterson D, Connelly RR, et al.** Clinical, epidemiologic, and virologic studies in four clusters of the chronic fatigue syndrome. *Arch Intern Med.* 1992;152:1611-6.
 62. **Mittenberg W, DiGiulio DV, Perrin S, Bass AE.** Symptoms following mild head injury: expectation as aetiology. *J Neurol Neurosurg Psychiatry.* 1992;55:200-4.
 63. **Schrader H, Obelieniene D, Bovim G, Surkiene D, Mickeviciene D, Misceviene I, et al.** Natural evolution of late whiplash syndrome outside the medicolegal context. *Lancet.* 1996;347:1207-11.
 64. **Wessely S.** Old wine in new bottles: neurasthenia and "ME." *Psychol Med.* 1990;20:35-53.
 65. **Mayou R, Radanov BP.** Whiplash neck injury. *J Psychosom Res.* 1996;40:461-74.
 66. **Karlsborg M, Smed A, Jespersen H, Stephensen S, Cortsen M, Jennum P, et al.** A prospective study of 39 patients with whiplash injury. *Acta Neurol Scand.* 1997;95:65-72.
 67. **Suarez FL, Savaiano DA, Levitt MD.** A comparison of symptoms after the consumption of milk or lactose-hydrolyzed milk by people with self-reported severe lactose intolerance. *N Engl J Med.* 1995;333:1-4.
 68. **Institute of Medicine.** Health Consequences of Service during the Persian Gulf War: Recommendations for Research and Information Systems. Washington, DC: National Academy Pr; 1996.
 69. **Evans RW.** Some observations on whiplash injuries. *Neurol Clin.* 1992;10:975-97.
 70. **Pennie B, Agambar L.** Patterns of injury and recovery in whiplash. *Injury.* 1991;22:57-9.
 71. **Simon GE, Daniell W, Stockbridge H, Claypoole K, Rosenstock L.** Immunologic, psychological, and neuropsychological factors in multiple chemical sensitivity. A controlled study. *Ann Intern Med.* 1993;119:97-103.
 72. **Fiedler N, Kipen HM, DeLuca J, Kelly-McNeil K, Natelson B.** A controlled comparison of multiple chemical sensitivities and chronic fatigue syndrome. *Psychosom Med.* 1996;58:38-49.
 73. **Morrison JD.** Fatigue as a presenting complaint in family practice. *J Fam Pract.* 1980;10:795-801.
 74. **Simon GE, Katon WJ, Sparks PJ.** Allergic to life: psychological factors in environmental illness. *Am J Psychiatry.* 1990;147:901-6.
 75. **Epstein SA, Kay G, Clauw D, Heaton R, Klein D, Krupp L, et al.** Psychiatric disorders in patients with fibromyalgia. A multicenter investigation. *Psychosomatics.* 1999;40:57-63.
 76. **Aaron LA, Bradley LA, Alarcon GS, Alexander RW, Triana-Alexander MA, Martin MY, et al.** Psychiatric diagnoses in patients with fibromyalgia are related to health care-seeking behavior rather than to illness. *Arthritis Rheum.* 1996;39:436-45.
 77. **Goldenberg DL.** An overview of psychologic studies in fibromyalgia. *J Rheumatol.* 1989;16:12-4.
 78. **Goldenberg DL.** Psychological symptoms and psychiatric diagnosis in patients with fibromyalgia. *J Rheumatol Suppl.* 1989;19:127-30.
 79. **Merskey H.** Physical and psychological considerations in the classification of fibromyalgia. *J Rheumatol.* 1989;16:72-9.
 80. **Payne TC, Leavitt F, Garron DC, Katz RS, Golden HE, Glickman PB, et al.** Fibrositis and psychological disturbance. *Arthritis Rheum.* 1982;25:213-7.
 81. **Drossman DA, McKee DC, Sandler RS, Mitchell CM, Cramer EM, Lowman BC, et al.** Psychosocial factors in the irritable bowel syndrome. A multivariate study of patients and nonpatients with irritable bowel syndrome. *Gastroenterology.* 1988;95:701-8.
 82. **Mendeloff AI, Monk M, Siegel CI, Lilienfeld A.** Illness experience and life stresses in patients with irritable colon and with ulcerative colitis. An epidemiologic study of ulcerative colitis and regional enteritis in Baltimore, 1960-1964. *N Engl J Med.* 1970;282:14-7.
 83. **Drossman DA, Lowman BC.** Irritable bowel syndrome: epidemiology, diagnosis and treatment. *Clin Gastroenterol.* 1985;14:559-73.
 84. **Walker EA, Roy-Byrne PP, Katon WJ, Li L, Amos D, Jiranek G.** Psychiatric illness and irritable bowel syndrome: a comparison with inflammatory bowel disease. *Am J Psychiatry.* 1990;147:1656-61.
 85. **Stewart DE.** Environmental hypersensitivity disorder, total allergy, and 20th-century disease: a critical review. *Can Fam Physician.* 1987;33:405-10.
 86. **Spaulding WB.** The clinical analysis of fatigue. *Appl Ther.* 1964;6:911-5.
 87. **French MA.** The clinical significance of tiredness. *Can Med Assoc J.* 1960;82:665-71.
 88. **Sugarman JR, Berg AO.** Evaluation of fatigue in a family practice. *J Fam Pract.* 1984;19:643-7.
 89. **Katerndahl DA.** Fatigue of uncertain etiology. *Fam Med.* 1983;1:26-38.
 90. **Jerrett WA.** Lethargy in general practice. *Practitioner.* 1981;225:731-7.
 91. **Allen PN.** Differential diagnosis of weakness and fatigue. *N Engl J Med.* 1994;231:414-8.
 92. **Katon WJ, Buchwald DS, Simon GE, Russo JE, Mease PJ.** Psychiatric illness in patients with chronic fatigue and those with rheumatoid arthritis. *J Gen Intern Med.* 1991;6:277-85.
 93. **Wood GC, Bentall RP, Gopfert M, Edwards RH.** A comparative psychiatric assessment of patients with chronic fatigue syndrome and muscle disease. *Psychol Med.* 1991;21:619-28.
 94. **Hickie I, Lloyd A, Hadzi-Pavlovic D, Parker G, Bird K, Wakefield D.** Can the chronic fatigue syndrome be defined by distinct clinical features? *Psychol Med.* 1995;25:925-35.
 95. **Manu P, Lane TJ, Matthews DA.** The frequency of the chronic fatigue syndrome in patients with symptoms of persistent fatigue. *Ann Intern Med.* 1988;109:554-6.
 96. **Gold D, Bowden R, Sixbey J, Riggs R, Katon WJ, Ashley R, et al.** Chronic fatigue. A prospective clinical and virologic study. *JAMA.* 1990;264:48-53.
 97. **Katon W, Russo J.** Chronic fatigue syndrome criteria. A critique of the requirement for multiple physical complaints. *Arch Intern Med.* 1992;152:1604-9.

98. Whitehead WE, Winget C, Fedoravicius AS, Wooley S, Blackwell B. Learned illness behavior in patients with irritable bowel syndrome and peptic ulcer. *Dig Dis Sci*. 1982;27:202-8.
99. Gough HG. Doctors' estimates of the percentage of patients whose problems do not require medical attention. *Med Educ*. 1977;11:380-4.
100. Derogatis LR, Lipman RS, Rickels K, Uhlenhuth EH, Covi L. The Hopkins Symptom Checklist (HSLC): a self-report symptom inventory. *Behav Sci*. 1974;19:1-15.
101. Walker EA, Keegan D, Gardner G, Sullivan M, Bernstein D, Katon WJ. Psychosocial factors in fibromyalgia compared with rheumatoid arthritis: I. Psychiatric diagnosis and functional disability. *Psychosom Med*. 1997;59:565-71.
102. Hammaren M, Hugoson A. Clinical psychiatric assessment of patients with burning mouth syndrome resisting oral treatment. *Swed Dent J*. 1989;13:77-88.
103. Yontchev E, Carlsson GE. Long-term follow-up of patients with orofacial discomfort complaints. *J Oral Rehabil*. 1992;19:13-9.
104. Miller MH, Topliss DJ. Chronic upper limb pain syndrome (repetitive strain injury) in the Australian workforce: a systematic cross sectional rheumatological study of 229 patients. *J Rheumatol*. 1988;15:1705-12.
105. Crisp AH, Moldofsky H. A psychosomatic study of writers' cramp. *Br J Psychiatry*. 1965;111:841-58.
106. Lucire Y. Neurosis in the workplace. *Med J Aust*. 1986;145:323-7.
107. Barsky AJ. Amplification, somatization, and the somatoform disorders. *Psychosomatics*. 1992;33:28-34.
108. Barsky AJ, Borus JF. Somatization and medicalization in the era of managed care. *JAMA*. 1995;274:1931-4.
109. Pennebaker JW. *The Psychology of Physical Symptoms*. New York: Springer-Verlag; 1982.
110. Barsky AJ, Wyshak G. Hypochondriasis and somatosensory amplification. *Br J Psychiatry*. 1990;157:404-9.
111. Ford CV. *The Somatizing Disorders: Illness as a Way of Life*. New York: Elsevier Biomedical; 1983.
112. Stuart S, Noyes R Jr. Attachment and interpersonal communication in somatization. *Psychosomatics*. 1999;40:34-43.
113. Pennebaker JW. Psychological bases of symptom reporting: perceptual and emotional aspects of chemical sensitivity. *Toxicol Ind Health*. 1994;10:497-511.
114. Pennebaker JW, Epstein D. Implicit psychophysiology: effects of common beliefs and idiosyncratic physiological responses on symptom reporting. *J Pers*. 1983;51:468-96.
115. Pennebaker JW, Watson D. The psychology of somatic symptoms. In: Kirmayer LJ, Robbins JM, eds. *Current Concepts of Somatization: Research and Clinical Perspectives*. Washington, DC: American Psychiatric Pr; 1991:24-35.
116. Alderman MH, Lampert B. Labeling of hypertensives: a review of the data. *J Clin Epidemiol*. 1990;43:195-200.
117. Haynes RB, Sackett DL, Taylor DW, Gibson ES, Johnson AL. Increased absenteeism from work after detection and labeling of hypertensive patients. *N Engl J Med*. 1978;299:741-4.
118. Dworkin RH, Harkstein G, Rosner HL, Walther RR, Sweeney EW, Brand L. A high-risk method for studying psychosocial antecedents of chronic pain: the prospective investigation of herpes zoster. *J Abnorm Psychol*. 1992;101:200-5.
119. Wielgosz AT, Earp J. Perceived vulnerability to serious heart disease and persistent pain in patients with minimal or no coronary disease. *Psychosom Med*. 1986;48:118-24.
120. Cope H, David A, Pelosi A, Mann A. Predictors of chronic "postviral" fatigue. *Lancet*. 1994;344:864-8.
121. Lees-Haley PR, Brown RS. Biases in perception and reporting following a perceived toxic exposure. *Percept Mot Skills*. 1992;75:531-44.
122. Marshall PD, O'Connor M, Hodgkinson JP. The perceived relationship between neck symptoms and precedent injury. *Injury*. 1995;26:17-9.
123. McFarland C, Ross M, DeCourville N. Women's theories of menstruation and biases in recall of menstrual symptoms. *J Pers Soc Psychol*. 1989;57:522-31.
124. Croyle RT, Sande GN. Denial and confirmatory search: paradoxical consequences of medical diagnosis. *J Appl Soc Psychol*. 1988;18:473-90.
125. Elks ML. On the genesis of somatization disorder: the role of the medical profession. *Med Hypotheses*. 1994;43:151-4.
126. Self-reported illness and health status among Gulf War veterans. A population-based study. The Iowa Persian Gulf Study Group. *JAMA*. 1997;277:238-45.
127. Myers MG, Cairns JA, Singer J. The consent form as a possible cause of side effects. *Clin Pharmacol Ther*. 1987;42:250-3.
128. Petrie K, Moss-Morris R, Weinman J. The impact of catastrophic beliefs on functioning in chronic fatigue syndrome. *J Psychosom Res*. 1995;39:31-7.
129. Radanov BP, Sturzenegger M, Di Stefano G. Long-term outcome after whiplash injury. A 2-year follow-up considering features of injury mechanism and somatic, radiologic, and psychosocial findings. *Medicine (Baltimore)*. 1995;74:281-97.
130. Schweiger A, Parducci A. Nocebo: the psychological induction of pain. *Pavlov J Biol Sci*. 1981;16:140-3.
131. Baker DB, Greenland S, Mendlein J, Harmon P. A health study of two communities near the Stringfellow Waste Disposal site. *Arch Environ Health*. 1988;43:325-34.
132. David AS, Wessely SC. The legend of Camelford: medical consequences of a water pollution accident [Editorial]. *J Psychosom Res*. 1995;39:1-9.
133. Roht LH, Vernon SW, Weir FW, Pier SM, Sullivan P, Reed LJ. Community exposure to hazardous waste disposal sites: assessing reporting bias. *Am J Epidemiol*. 1985;122:418-33.
134. Kirmayer LJ, Taillefer S. Somatoform disorders. In: Turner SM, Hersen M, eds. *Adult Psychopathology and Diagnosis*. 3d ed. New York: J Wiley; 1997:333-82.
135. McDaniel SH, Hepworth J, Doherty WJ. *Medical Family Therapy: A Biopsychosocial Approach to Families with Health Problems*. New York: Basic Books; 1992.
136. Block AR, Kremer EF, Gaylor M. Behavioral treatment of chronic pain: the spouse as a discriminative cue for pain behavior. *Pain*. 1980;9:243-52.
137. Jamison RN, Virts KL. The influence of family support on chronic pain. *Behav Res Ther*. 1990;28:283-7.
138. Craig TK, Drake H, Mills K, Boardman AP. The South London Somatization Study. II. Influence of stressful life events, and secondary gain. *Br J Psychiatry*. 1994;165:248-58.
139. Waxler NE. The social labeling perspective on illness and medical practice. In: Eisenberg L, Kleinman A, eds. *The Relevance of Social Science for Medicine*. Boston: Reidel; 1980:283-306.
140. Goldenberg DL, Mossey CJ, Schmid CH. A model to assess severity and impact of fibromyalgia. *J Rheumatol*. 1995;22:2133-8.
141. Satel SL. When disability benefits make patients sicker [Editorial]. *N Engl J Med*. 1995;333:794-6.
142. Bellamy R. Compensation neurosis: financial reward for illness as nocebo. *Clin Orthop*. 1997;336:94-106.
143. Robertson LS, Keeve JP. Worker injuries: the effects of Workers' Compensation and OSHA inspections. *J Health Polit Policy Law*. 1983;8:581-97.
144. Higgs PE, Edwards D, Martin DS, Weeks PM. Carpal tunnel surgery outcomes in workers: effect of workers' compensation status. *J Hand Surg [Am]*. 1995;20:354-60.
145. Hadler NM. If you have to prove you are ill, you can't get well. *Spine*. 1996;21:2397-400.
146. Ireland DC. Repetition strain injury: the Australian experience—1992 update. *J Hand Surg [Am]*. 1995;20(3 Pt 2):S53-6.
147. Rahe RH, Arthur RJ. Life change and illness studies: past history and future directions. *J Human Stress*. 1978;4:3-15.
148. Tessler R, Mechanic D. Psychological distress and perceived health status. *J Health Soc Behav*. 1978;19:254-62.
149. Robbins PR, Meyersburg HA, Tanck RH. Interpersonal stress and physical complaints. *J Pers Assess*. 1974;38:578-85.
150. Drossman DA. Patients with psychogenic abdominal pain: six years' observation in the medical setting. *Am J Psychiatry*. 1982;139:1549-57.
151. Kellner R. *Somatization and Hypochondriasis*. New York: Praeger; 1986.
152. Mechanic D. Effects of psychological distress on perceptions of physical health and use of medical and psychiatric facilities. *J Human Stress*. 1978;4:26-32.
153. Ryan GA, Bampton M. Comparison of data process operators with and without upper limb symptoms. *Community Health Stud*. 1988;12:63-8.
154. Linton SJ, Kamwendo K. Risk factors in the psychosocial work environment for neck and shoulder pain in secretaries. *J Occup Med*. 1989;31:609-13.
155. Hadler NM. Back pain in the workplace. What you lift or how you lift matters far less than whether you lift or when [Editorial]. *Spine*. 1997;22:935-40.
156. Magora A. Investigation of the relation between low back pain and occupation. V. Psychological aspects. *Scand J Rehabil Med*. 1973;5:191-6.
157. Papageorgiou AC, Macfarlane GJ, Thomas E, Croft PR, Jayson MI, Silman AJ. Psychosocial factors in the workplace—do they predict new episodes of low back pain? Evidence from the South Manchester Back Pain Study. *Spine*. 1997;22:1137-42.
158. Hadler NM. Workers with disabling back pain [Editorial]. *N Engl J Med*. 1997;337:341-3.
159. Hall EM, Johnson JV. A case study of stress and mass psychogenic illness in industrial workers. *J Occup Med*. 1989;31:243-50.
160. Affleck G, Tennen H, Urrows S, Higgins P. Person and contextual features of daily stress reactivity: individual differences in relations of undesirable daily events with mood disturbance and chronic pain intensity. *J Pers Soc Psychol*. 1994;66:329-40.
161. Anderson KO, Bradley LA, Young LD, McDaniel LK, Wise CM. Rheumatoid arthritis: review of psychological factors related to etiology, effects, and treatment. *Psychol Bull*. 1985;98:358-87.
162. Smed A. Cognitive function and distress after common whiplash injury. *Acta Neurol Scand*. 1997;95:73-80.
163. Escobar JI, Canino G, Rubio-Stipec M, Bravo M. Somatic symptoms after a natural disaster: a prospective study. *Am J Psychiatry*. 1992;149:965-7.
164. McFarlane AC, Atchison M, Rafalowicz E, Papay P. Physical symptoms in post-traumatic stress disorder. *J Psychosom Res*. 1994;38:715-26.
165. Bravo M, Rubio-Stipec M, Canino GJ, Woodbury MA, Ribera JC. The psychological sequelae of disaster stress prospectively and retrospectively evaluated. *Am J Community Psychol*. 1990;18:661-80.
166. Solomon Z. Somatic complaints, stress reaction, and posttraumatic stress disorder: a three-year follow-up study. *Behav Med*. 1988;14:179-85.
167. Shalev A, Bleich A, Ursano RJ. Posttraumatic stress disorder: somatic comorbidity and effort tolerance. *Psychosomatics*. 1990;31:197-203.
168. Koss MP, Koss PG, Woodruff WJ. Deleterious effects of criminal victimization on women's health and medical utilization. *Arch Intern Med*. 1991;151:342-7.
169. Waylonis GW, Perkins RH. Post-traumatic fibromyalgia. A long-term follow-up. *Am J Phys Med Rehabil*. 1994;15:541-8.
170. Hazlett RL, Haynes SN. Fibromyalgia: a time-series analysis of the stressor-physical symptom association. *J Behav Med*. 1992;15:541-58.
171. Dailey PA, Bishop GD, Russell IJ, Fletcher EM. Psychological stress and the fibrositis/fibromyalgia syndrome. *J Rheumatol*. 1990;17:1380-5.

172. **Beebe GW.** Follow-up studies of World War II and Korean War prisoners. II. Morbidity, disability, and maladjustments. *Am J Epidemiol.* 1975;101:400-22.
173. **Sternbach RA.** Psychological dimensions and perceptual analyses, including pathologies of pain. In: Carterett ED, Friedman MD, eds. *Handbook of Perception.* New York: Academic Pr; 1978:231-61.
174. **Salovey P, Birnbaum D.** Influence of mood on health-relevant cognitions. *J Pers Soc Psychol.* 1989;57:539-51.
175. **Larsen RJ.** Neuroticism and selective encoding and recall of symptoms: evidence from a combined concurrent-retrospective study. *J Pers Soc Psychol.* 1992;62:480-8.
176. **Kirmayer LJ, Robbins JM, Paris J.** Somatoform disorders: personality and the social matrix of somatic distress. *J Abnorm Psychol.* 1994;103:125-36.
177. **Imboden JB, Canter A, Cluff LE.** Convalescence from influenza. A study of the psychological and clinical determinants. *Arch Intern Med.* 1961;108:393-9.
178. **Lucock MP, Morley S, White C, Peake MD.** Responses of consecutive patients to reassurance after gastroscopy: results of a self-administered questionnaire survey. *BMJ.* 1997;315:572-5.
179. **Potts SG, Bass CM.** Psychosocial outcome and use of medical resources in patients with chest pain and normal or near-normal coronary arteries: a long-term follow-up study. *Q J Med.* 1993;86:583-93.
180. **Weber BE, Kapoor WN.** Evaluation and outcomes of patients with palpitations. *Am J Med.* 1996;100:138-48.
181. **Twemlow SW, Bradshaw SL Jr, Coyne L, Lerma BH.** Patterns of utilization of medical care and perceptions of the relationship between doctor and patient with chronic illness including chronic fatigue syndrome. *Psychol Rep.* 1997;80:643-58.
182. **Russo J, Katon W, Sullivan M, Clark M, Buchwald D.** Severity of somatization and its relationship to psychiatric disorders and personality. *Psychosomatics.* 1994;35:546-56.
183. **Kroenke K, Spitzer RL, Williams JB, Linzer M, Hahn SR, deGruy FV 3d, et al.** Physical symptoms in primary care. Predictors of psychiatric disorders and functional impairment. *Arch Fam Med.* 1994;3:774-9.
184. **Kisely S, Goldberg D, Simon G.** A comparison between somatic symptoms with and without clear organic cause: results of an international study. *Psychol Med.* 1997;27:1011-9.
185. **Dworkin SF, Von Korff M, LeResche L.** Multiple pains and psychiatric disturbance. An epidemiologic investigation. *Arch Gen Psychiatry.* 1990;47:239-44.
186. **Clark DM, Salkovskis PM, Hackmann A, Wells A, Fennell M, Ludgate J, et al.** Two psychological treatments for hypochondriasis. A randomized controlled trial. *Br J Psychiatry.* 1998;173:218-25.
187. **Salkovskis PM.** Somatic problems. In: Hawton K, Salkovskis PM, Kirk JW, Clark DM, eds. *Cognitive Behaviour Therapy for Psychiatric Problems: A Practical Guide.* New York: Oxford Univ Pr; 1989:235-76.
188. **Skinner JB, Erskine A, Pearce S, Rubinstein I, Taylor M, Foster C.** The evaluation of a cognitive behavioural treatment programme in outpatients with chronic pain. *J Psychosom Res.* 1990;34:13-20.
189. **Speckens AE, van Hemert AM, Spinhoven P, Hawton KE, Bolk JH, Rooijmans HG.** Cognitive behavioural therapy for medically unexplained physical symptoms: a randomised controlled trial. *BMJ.* 1995;311:1328-32.
190. **DeGuire S, Gevirtz R, Kawahara Y, Maguire W.** Hyperventilation syndrome and the assessment of treatment for functional cardiac symptoms. *Am J Cardiol.* 1992;70:673-7.
191. **van Dulmen AM, Fennis JF, Bleijenberg G.** Cognitive-behavioral group therapy for irritable bowel syndrome: effects and long-term follow-up. *Psychosom Med.* 1996;58:508-14.
192. **Deale A, Chalder T, Marks I, Wessely S.** Cognitive behavior therapy for chronic fatigue syndrome: a randomized controlled trial. *Am J Psychiatry.* 1997;154:408-14.
193. **Blanchard EB, Appelbaum KA, Radnitz CL, Michultka D, Morrill B, Kirsch C, et al.** Placebo-controlled evaluation of abbreviated progressive muscle relaxation and of relaxation combined with cognitive therapy in the treatment of tension headache. *J Consult Clin Psychol.* 1990;58:210-5.
194. **Buckelew SP.** Fibromyalgia: a rehabilitation approach. A review. *Am J Phys Med Rehabil.* 1989;68:37-42.
195. **Keefe FJ, Dunsmore J, Burnett R.** Behavioral and cognitive-behavioral approaches to chronic pain: recent advances and future directions. *J Consult Clin Psychol.* 1992;60:528-36.
196. **Martin PR, Nathan PR, Milech D, van Keppel M.** Cognitive therapy vs. self-management training in the treatment of chronic headaches. *Br J Clin Psychol.* 1989;28:347-61.
197. **Sharpe M.** Cognitive behavioural therapies in the treatment of functional somatic symptoms. In: Mayou R, Bass C, Sharpe M, eds. *Treatment of Functional Somatic Symptoms.* New York: Oxford Univ Pr; 1995:122-43.
198. **Mayou RA, Bryant BM, Sanders D, Bass C, Klimes I, Forfar C.** A controlled trial of cognitive behavioural therapy for non-cardiac chest pain. *Psychol Med.* 1997;27:1021-31.
199. **Sharpe M, Peveler R, Mayou R.** The psychological treatment of patients with functional somatic symptoms: a practical guide. *J Psychosom Res.* 1992;36:515-29.
200. **Hellman CJ, Budd M, Borysenko J, McClelland DC, Benson H.** A study of the effectiveness of two group behavioral medicine interventions for patients with psychosomatic complaints. *Behav Med.* 1990;16:165-73.
201. **Sharpe M, Hawton K, Simkin S, Surawy C, Hackmann A, Klimes I, et al.** Cognitive behaviour therapy for the chronic fatigue syndrome: a randomized controlled trial. *BMJ.* 1996;312:22-6.
202. **Payne A, Blanchard EB.** A controlled comparison of cognitive therapy and self-help support groups in the treatment of irritable bowel syndrome. *J Consult Clin Psychol.* 1995;63:779-86.
203. **Fulcher KY, White PD.** Randomised controlled trial of graded exercise in patients with the chronic fatigue syndrome. *BMJ.* 1997;314:1647-52.
204. **Lidbeck J.** Group therapy for somatization disorders in general practice: effectiveness of a short cognitive-behavioural treatment model. *Acta Psychiatr Scand.* 1997;96:14-24.